


The Distribution of Dark Matter in the Universe on scales of $10^{10} M_{\odot}$ to $10^{15} M_{\odot}$ ^a

Luís Teodoro and Michael S. Warren ^b
Theoretical Division, Los Alamos National Laboratory
Los Alamos, New Mexico, USA



The use of parallel computers and increasingly sophisticated software has allowed us to perform a large suite of N-body simulations using from 10^8 to 10^9 particles. We will report on our recent convergence tests of the halo mass function, N-point correlation functions, power spectrum and pairwise velocity from very large high resolution treecode N-body simulations. Rather than basing results on just one or two large simulations, now one can investigate the role of different numerical and physical effects on the statistics used to characterize the mass distribution of the Universe.

1 Introduction

Over the last two decades cosmological N -body simulations have played a crucial role in the study of the formation and evolution of cosmic structure. In this approach, the initial statistical properties of the fluctuations, at some early stage of the universe, are set using linear theory. N -body simulations are then used to evolve the structure into the deeply nonlinear regime. This dynamical state is then compared with the large-scale structure in the galaxy data-sets.

^aTalk presented by L. Teodoro at the XXXIXth Rencontres de Moriond on *Exploring the Universe*, La Thuile, Italy, March 28 - April 4, 2004.

^bteodoro@lanl.gov(LT), msw@lanl.gov(MSW)

The impressive progress achieved in the observational front with the completion of very large surveys such as 2dF and SDSS poses a clear challenge to the numerical work in cosmology: the precision of the predictions provided by the current N -body experiments have to be of the order of a few percent.

2 Numerical Simulations

The cosmological model adopted is a low-density, flat, Λ CDM Universe with the parameters $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$ and $h = 0.7$. The power spectrum of the initial conditions was set up using the output transfer function of CMBFAST¹, assuming $\Omega_b = 0.04$ and a normalisation of $\sigma_8 = 0.9$. The entire suite of simulations were performed using the Hashed Oct-Tree code (HOT)², a parallel tree code with periodic boundary conditions. The simulation parameters are listed in Table 1.

Table 1: Suite of simulations

Run	Ω_m	Ω_Λ	h	# Part.	L_{box} [Mpc h^{-1}]	ϵ [kpc h^{-1}]	m_p [$M_\odot$]
dtd10	0.30	0.70	0.70	512 ³	1536	98.0	2.3×10^{11}
dtd11	0.30	0.70	0.70	512 ³	768	49.0	2.8×10^{10}
dtd12	0.30	0.70	0.70	512 ³	384	24.5	3.5×10^9
ej1	0.30	0.70	0.70	512 ³	192	12.3	4.4×10^9
ef10	0.30	0.70	0.70	512 ³	96	4.9	5.5×10^8
ef4	0.30	0.70	0.70	768 ³	1152	65.5	2.8×10^{11}
ef7	0.30	0.70	0.70	768 ³	288	14.0	4.4×10^9
ei6	0.30	0.70	0.70	1024 ³	768	24.5	3.5×10^9

3 Analysis

The main purpose of this analysis is to understand how changes in mass resolution, particle number and box size affect some well known estimators applied in Large Scale Structure studies. For each cosmological volume we estimate the two-point correlation function, the power spectra, the mass function and the mean pairwise velocity, see Figure 1. The color map is shown in Table 1.

The two-point correlation function is presented in Figure 1 (a). On small scales, the amplitude of the two-point correlation function is suppressed by softening gravity and lack of small scale power in the initial conditions. This affects the estimates in a range which roughly extends up to five times the size of the smoothing kernel [see arrows in top left panel of the abovementioned figure]. The flattening of the two-point correlation function seems to disappear once the mass resolution (softening) increases (decreases). On scales where $\xi \approx 1$ all the simulations show a rather good agreement among themselves. The box size only looks to matter on scales well within the linear regime ($\xi \ll 1$) where the smaller boxes present a slightly smaller amplitude. The solid (dashed) lines represent the two-point Smith *et al*³ (linear) correlation function.

To measure the power spectra of our simulations we applied the technique detailed in Jenkins *et al*⁴ and Smith *et al*³. In doing so discreteness and grid effects are taken in account and removed from the final estimation. As expected, the power spectra of the different N -body simulations show the same features as the two-point correlation function. The analytical expression for the non-linear power spectra due to Smith *et al*³ and the linear power spectra are shown by the solid and dashed black lines, respectively. Further details can be found in Warren & Teodoro⁵.

We used the friends-of-friends algorithm⁶ to identify dark matter halos. This halo finder depends on just one parameter, b , which defines the linking length as $bn^{-1/3}$ where n is the mean particle density; we follow the prescription of Jenkins *et al*⁷ and set $b = 0.2$. Figure (c) demonstrates that the agreement between the analytical expressions of Jenkins *et al*⁷ (solid line) and Sheth and Tormen⁸ (dotted line) and our estimates of the mass function is excellent. The small departures seen in the mass range $M > 10^{14}h^{-1}M_{\odot}$ are mainly caused by small number statistics.

Finally, Figure 1 (d) shows the mean pairwise velocity, $v_{12}(r)$, as function of pair separation. The Hubble line, given by $v_{Hubble} = -Hr$, is also shown. The mean pairwise velocity vanishes at the smallest separations resolved in our simulations. In the subset of simulations which better probe small scales (small volume and larger mass resolution) $v_{12}(r)$ closely follows the Hubble line up to $\sim 300 \text{ kpc } h^{-1}$. This indicates that structures on such scales are close to relaxation. The mean pairwise velocity reaches a minimum in scales comparable to the correlation length [see Figure 1(d)]. At larger scales the $v_{12}(r)$ intersects the Hubble line and, at very large separations it decays to zero, in accordance with the principle of large-scale homogeneity and isotropy.

Some simulations in this suite have been further analyze in Seljak & Warren⁹.

Acknowledgments

We thank A.R. Jenkins, G. Evrard, K. Abazajian and S. Habib for help and useful discussions. This work was performed under the auspices of the U.S. Dept. of Energy, and supported by its contract #W-7405-ENG-36 to Los Alamos National Laboratory. Simulations were performed on the Space Simulator Beowulf cluster at Los Alamos National Laboratory.

References

1. U. Seljak & M. Zaldarriaga, *ApJ*. **469**, 437 (1996)
2. M.S. Warren & J.K. Salmon, *Supercomputing '93*, IEEE Comp. Soc., Los Alamitos 1993
3. R. Smith, J.A. Peacock, A. Jenkins, S.D.M. White, C.S. Frenk, F.R. Pearce, P.A. Thomas, G. Efstathiou, H.M.P. Couchman, *MNRAS* **341**, 1311 (2003)
4. A. Jenkins, C.F. Frenk, F.P. Pearce, P.A. Thomas, J.M. Colberg, S.D.M. White, H.M.P. Couchman, J.A. Peacock, G. Efstathiou, A.H. Nelson, *ApJ*. **499**, 20 (1998)
5. M.S. Warren & L. Teodoro, in preparation.
6. M. Davis, G. Efstathiou, C.S. Frenk and S.D.M. White, *ApJ*. **292**, 371 (1985); D. Pfitzner, J. Salmon and T. Sterling, *Journal of Data Mining and Knowledge Discovery* **1**, 419 (1977)
7. A. Jenkins, C.S. Frenk, S.D.M. White, J.M. Colberg, S. Cole, A.E. Evrard, H.M.P. Couchman, N. Yoshida, *MNRAS* **321**, 372 (2001)
8. R. Sheth & G. Tormen, *MNRAS* **308**, 119 (1999)
9. U. Seljak & M.S. Warren, ArXiv e-print astro-ph/0403698
10. W.H. Press & P. Schechter, *ApJ*. **187**, 425 (1974)




Figure 1: (a) Two point correlation function, (b) power spectrum, (c) mass function and (d) mean pairwise velocity. See color map in Table 1. The arrows in the top left panel indicate the gravitational softening. In the same panel, we also show the linear theory (dashed line) and the Smith *et al* (solid line) two-point correlation function fitting for the assumed cosmological model. The solid, dotted and dashed lines in panel (c) show the Sheth-Tormen, Jenkins *et al* and Press-Schechter¹⁰ mass functions, respectively. Finally, in panel (d) the dashed line represents the Hubble flow.